



## Lec. 1

# Examination of the Mouth and Other Relevant Structures

Successful pediatric dental care requires a detailed history, thorough examination, investigations, accurate diagnosis, and a well-planned treatment strategy. Dentists must collect all relevant patient and family information, obtain informed consent, and explain the treatment plan to parents. This includes not only corrective treatment but also a prevention plan to establish a long-term “dental home.” The first dental visit should occur with the eruption of the first tooth or by 12 months of age, as early detection and management improve both oral and general health of the child.

**Obtaining accurate data in a child is very difficult. The reasons may be any of the following:**

- 1) Most of the times, it is the parent or the guardian who will be providing the required data about the child and not the child himself or herself.
- 2) It is impossible to observe everything a child does or says and make accurate records of what goes on.
- 3) Most children do not behave in the dental clinic the same way as they do at home or with their friends or teachers.
- 4) Data reported by parents and teacher may be inaccurate.
- 5) Information provided by parents or guardians is dependable on their emotional maturity.
- 6) Unless reports are made immediately after the observation, the parents may forget to mention minor yet important findings.
- 7) There may be a deliberate distortion by the observer to show the child or the parent in a favorable light
  - The history facilitates the diagnosis of many conditions, even before the hands-on examination.
  - Because there are often specific questions pertinent to a child's medical history that will be relevant to their management, it is desirable that parents be present.



- The successful practice of pediatric dentistry is not merely the completion of any operative procedure but also ensuring a positive dental outcome for the future oral health behavior of that individual and family.

### **Treatment planning:**

Treatment planning is the step-by-step organization of a child's dental needs to ensure maximum benefit. It should address current problems, prevent future issues, and be based on a complete history and examination. Parental consent is required, and discussions with parents—known as anticipatory guidance—help implement the plan effectively.

### **Advantages of Treatment Planning**

- 1) Avoiding the re-diagnosis at every visit.
- 2) Give serial appointments on the first day as the patient's treatment needs that are already planned in a sequential order (step-by-step guideline).
- 3) Instruments can be prepared well in advance before the patient's arrival for the treatment.
- 4) Estimation of the time and no. of appointments required as well as the total fee.

⊕ Treatment plan must be discussed with the parents and permission taken before performing any treatment on the child.

### **The followings information must be taken:**

- ✓ Dental need of their child including the treatment as well as the preventive measures.
- ✓ Amount of time required to perform the treatment and the total cost.

### **Treatment planning may be modified during the procedure based on the following:**

- 1) Estimation of cooperation from the patient and parents.
- 2) Assessment of the condition of the teeth and the oral hygiene.
- 3) Whether extraction is needed or not.
- 4) Nature of tooth movement and type of appliance required.



## THE DIAGNOSTIC METHOD

Successful pediatric dental care begins with an accurate diagnostic process. This requires a comprehensive history, thorough clinical and radiographic examination, and careful interpretation of findings to reach a precise diagnosis and guide effective treatment planning.

Diagnosis and treatment planning involve gathering all relevant facts—from patient and parental reports, clinical signs, and diagnostic aids—and analyzing them systematically to determine both immediate needs and long-term care strategies.

Before making a diagnosis and developing a treatment plan, the dentist must collect and evaluate the facts associated with the patient's or parents' chief concern and any other identified problems that may be unknown to the patient or parents. Some pathognomonic signs may lead to an almost immediate diagnosis. For example, obvious gingival swelling and drainage may be associated with a single, badly carious primary molar. Although these associated facts are collected and evaluated rapidly, they provide a diagnosis only for a single problem area. On the other hand, sometimes there is a need to postpone a comprehensive diagnosis of all of the patient's problems or potential problems until conditions that are more urgent are resolved. For example, a patient with necrotizing ulcerative gingivitis (NUG) or a newly fractured crown needs immediate treatment, but the treatment will likely be only palliative, and further diagnostic and treatment procedures will be required later.

**A thorough examination of the pediatric dental patient includes an assessment of the following:**

- General growth and health
- Chief complaint, such as pain
- Extraoral soft tissue and TMJ evaluation
- Intraoral soft tissue
- Oral hygiene and periodontal health
- Intraoral hard tissue
- Developing occlusion
- Caries risk
- Behavior



Additional diagnostic aids are often also required, such as radiographs, study models, photographs, pulp tests, and, infrequently, laboratory tests.

### Components of oral examination and diagnosis

- Recording the history
- Examination of the patient
- Provisional diagnosis
- Special examination
- Final diagnosis
- Treatment plan (including medical referrals).

**1-Recording the history:** Recording the history (medical, dental and familial). So, to get a correct diagnosis, gathering complete and comprehensive Information about the followings should be done:

- **Vital statistics** (demographic information about the child like: birth, recognition, and social structure).
- **Chief complaint** and History of present illness (This is concerned about what made the patient to visit the dentist or what they are seeking from treatment).
- **Family (social) history** and pre- and postnatal history about the child (provides relevant information about the social background of the child and his family).
- **Medical and drug history.** It may affect the child's growth and development (social and psychological) as well as the type of diet taken by the child.

(Various diseases or functional disturbances may directly or indirectly cause or predispose to oral problems and may affect the delivery of oral care).

Information regarding the child's social and psychological development is important. Accurate information reflecting a child's learning, behavioral, or communication problems is sometimes difficult to obtain initially, especially when the parents are aware of their child's developmental disorder but are reluctant to discuss it. Mentally retarded child can be determined by asking question about the learning process and the child's behavioral and communication problem.



- **Drug history** (Details of the drugs being used for systemic ailments, any adverse reaction to drugs, and any drugs already used for the condition)
- **Behavioral history** (Include any behavior problems in the dental office, which are often related to the child's inability to communicate with the dentist and to follow instructions may call upon the need for behavior management or shaping).
- **Diet history** (Type of meal, Habits of snacking between meals should be evaluated as they may be cariogenic. In case of high cariogenic patients, a diet diary with number of sugar exposures should be noted while taking diet history).
- **Past dental history** (Previous care (treatment) in the dental office, Oral hygiene, habits, Previous and current fluoride therapy).

## 2- CLINICAL EXAMINATION

**Includes:** General Examination, Extra - and Intraoral examination.

### General Examination

General well-being of the child can be examined by a brief survey of the entire child's body. In addition to examining the oral cavity structures, the dentist may in some cases wish to note the patient's size, stature, gait, or involuntary movements. The first clue to malnutrition may come from observing a patient's abnormal size or stature. Similarly, the severity of a child's illness, even if oral in origin, may be recognized by observing a weak, unsteady gait or lethargy and malaise as the patient walks into the office.

**General examination includes the examination of the followings:**

- 1) Head, hair, face, neck and hands
- 2) Temporomandibular joint
- 3) Lymph nodes
- 4) Lips, Chin and tongue



## 1) Head, hair, face, neck and hands

The dentist can gather useful information while being acquainted with a new patient. After the patient seated in the dental chair, attention to the patient's hair, head, face, neck, and hands should be among the first observations made by the dentist, because abnormalities in size, shape, symmetry, function of head and neck indicate various syndromes that may be associated with oral abnormalities.

Inspection and palpation of the patient's head and neck are indicated. Unusual characteristics of the hair or skin should be noted. The dentist may observe signs of problems such as head lice, ringworm, or impetigo during the examination. Proper referral is indicated immediately, because these conditions are contagious.



Ringworm



Impetigo

**The patient's hands may reveal information pertinent to a comprehensive diagnosis (how?):**

- 1) The dentist may first detect an elevated temperature by holding the patient's hand.
- 2) Cold, clammy hands or bitten fingernails may be the first indication of abnormal anxiety in the child.
- 3) A callused or unusually clean digit suggests a persistent sucking habit.
- 4) Clubbing of the fingers or a bluish color in the nail beds suggests congenital heart disease, which may require special precautions during dental treatment.



## 2) Temporomandibular joint evaluation

The dentist should evaluate temporomandibular joint (TMJ) function and the associated musculature by palpating the head of each mandibular condyle and by observing the patient while the mouth is closed (teeth clenched), at rest, and in various open positions.

Movements of the condyles or jaw that do not flow smoothly or that deviate from the expected norm should be noted. Similarly, any crepitus (that may be heard or identified by palpation) as well as any other abnormal sounds or clicking should be noted. Sore masticatory muscles (that may be detected by pain, deviation and restricted mouth opening) may also signal TMJ dysfunction. Such deviations from normal TMJ function may require further evaluation and treatment. There is a consensus that temporomandibular disorders in children can be managed effectively by the following conservative and reversible therapies:

patient education, mild physical therapy, behavioral therapy, medications, and occlusal splints

Mouth opening (which is normally 40–45 mm) is also related to TMJ function and should be examined.

## 3) Lymph nodes

The extra oral examination continues with palpation of the patient's neck and submandibular area and any deviations from normal, such as unusual tenderness or enlargement, should be noted and follow up tests performed or referrals made as indicated.



#### 4) Lips, Chin and tongue

If the child is old enough to talk, speech should be evaluated.

The positions of the tongue, lips, and perioral musculature during speech, while swallowing, and at rest may provide useful diagnostic Information.

- Chin prominence is related to mandibular position.
- Lips normally should be competent (touch each other lightly or with 0-1 mm of gap).
- When the lips do not approximate each other at rest, they are termed as incompetent.